

animal

<http://journals.cambridge.org/ANM>


Additional services for **animal**:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Prediction of beef eating quality in France using the Meat Standards Australia system

I. Legrand, J.-F. Hocquette, R. J. Polkinghorne and D. W. Pethick

animal / *FirstView* Article / August 2012, pp 1 - 6

DOI: 10.1017/S1751731112001553, Published online: 10 August 2012

Link to this article: http://journals.cambridge.org/abstract_S1751731112001553

How to cite this article:

I. Legrand, J.-F. Hocquette, R. J. Polkinghorne and D. W. Pethick Prediction of beef eating quality in France using the Meat Standards Australia system. animal, Available on CJO 2012 doi:10.1017/S1751731112001553

Request Permissions : [Click here](#)

Prediction of beef eating quality in France using the Meat Standards Australia system

I. Legrand¹, J.-F. Hocquette^{2†}, R. J. Polkinghorne³ and D. W. Pethick⁴

¹Institut de l'Elevage, Service Qualité des Viandes, MRAL, 87060 Limoges Cedex 2, France; ²INRA, UR1213 Herbivores, 63122 Saint-Genès-Champanelle, France; ³431 Timor Road, Murrurundi, NSW 2338, Australia; ⁴School of Veterinary and Biomedical Science, Murdoch University, Murdoch, Western Australia 6150, Australia

(Received 20 October 2011; Accepted 12 June 2012)

An experiment was set up for (i) comparing Australian and French consumer preferences to beef and to (ii) quantify how well the Meat Standards Australia (MSA) grading model could predict the eating quality of beef in France. Six muscles from 18 Australian and 18 French cattle were tested as paired samples. In France, steaks were grilled 'medium' or 'rare', whereas in Australia 'medium' cooking was used. In total, 360 French consumers took part in the 'medium' cooking test, with each eating half Australian beef and half French beef and 180 French consumers tested the 'rare' beef. Consumers scored steaks for tenderness (tn), juiciness (ju), flavour liking (fl) and overall liking (ov). They also assigned a quality rating to each sample: 'unsatisfactory', 'satisfactory everyday quality' (3*), 'better than everyday quality' (4*) or 'premium quality' (5*). The prediction of the final ratings (3*, 4*, 5*) by the French consumers using the MSA-weighted eating quality score ($0.3 \text{ tn} + 0.1 \text{ ju} + 0.3 \text{ fl} + 0.3 \text{ ov}$) was over 70%, which is at least similar to the Australian experience. The boundaries between 'unsatisfactory', 3*, 4* and 5* were found to be ca. 38, 61 and 80, respectively. The differences between extreme classes are therefore slightly more important in France than in Australia. On average, even though it does not have predictive equations for bull meat, the mean predicted scores calculated by the MSA model deviated from observed values by a maximum of 5 points on a 0 to 100 scale except for the Australian oyster blade and the French topside, rump and outside (deviating by <15). Overall, the data indicate that it would be possible to manage a grading system in France as there is high agreement and consistency across consumers. The 'rare' and 'medium' results are also very similar, indicating that a common set of weightings and cut-offs can be employed.

Keywords: beef quality, palatability, prediction, consumers, sensory evaluation

Implications

Beef is characterized by a high variability of its palatability, which is one reason for consumer dissatisfaction. In France, there is still no reliable technique to predict beef quality to deliver a consistent eating experience for the consumer. Australia has developed the Meat Standards Australia (MSA) grading scheme to predict beef quality for each individual 'muscle × specific cooking method' combination using various information on the corresponding animals and meats. The MSA system has been shown to predict beef quality not only in Australia, but also in many other countries (Hwang *et al.*, 2008; Thompson *et al.*, 2010). The results of this study indicate that it would be possible to manage a grading system in France similar to the MSA system. In conclusion, an MSA-like meat grading system may be set up in France.

Introduction

Australia has developed the Meat Standards Australia (MSA) grading scheme to predict beef quality for consumers (Ferguson *et al.*, 1999; Polkinghorne *et al.*, 1999; Thompson *et al.*, 1999a and 1999b; Thompson, 2002). This system is based on the development and the use of a research database with a large amount of data, including the use of a large-scale consumer testing system with cuts cooked in different ways, as well as information on the corresponding animals, carcasses and meats. The system is also based on statistical analyses carried out on this database to identify the critical control points of beef palatability, which is indicated for individual muscles and for a specific cooking method and ageing time (Watson *et al.*, 2008b).

France is the country in the European Union (EU) with the largest cattle herd. France is therefore the largest beef meat producer and consumer in Europe, even though the

† E-mail: jfhocquette@clermont.inra.fr

Table 1 Some characteristics of the animal groups

	Australia	France	
	Steers	Young bulls	Cows
Number of animals	18	3	15
Number of muscles per animal	6	6	6
Number of degrees of doneness	1	2	2
Number of samples = number of animals × number of muscles × number of degrees of doneness	108	36	180
USDA ossification	141	150	457
USDA marbling	302	340	348
Carcass weight (kg)	319	282	329

USDA = United States Department of Agriculture.

consumption of beef has declined to reach 25 kg/person per year in 2010 (Hocquette and Chatellier, 2011).

French beef meat operators have been informed about the MSA programme (Moëvi *et al.*, 2008a and 2008b), which was (and is still) promoted at the international level. The French Livestock Institute ('Institut de l'Élevage', a non-profit national and technical institute dealing with applied research and development and specialized in livestock farming and products – cattle, sheep, goats, horses, forage resources and rural land use) and 'INRA de Theix' (public basic research institute for agriculture in France) were tasked to assess the MSA system with perspectives for the French beef industry. This work was also undertaken as part of the current European research programme ProSafeBeef (a programme focused on beef quality and safety, 2007 to 2011). The study was supported by the French Meat and Livestock Association (Interbev) and the National Office for Meat and Dairy Products (now FranceAgriMer). This work was carried out in 2007/2008, leading to a comprehensive report (Moëvi *et al.*, 2008a) and a viewpoint publication (Hocquette *et al.*, 2011a). As a result of this work, experts involved in the French meat industry recognized many qualities of the MSA system. It was judged comprehensive, consistent and scientifically supported and finally very interesting to be described and debated in France.

On the basis of these observations, an experiment described in this paper was set up to test the accuracy for predicting palatability scores of French consumers when using the MSA sensory protocol and prediction model. This paper reports the results of this experiment, which allowed the following comparisons to be made: (i) between French and Australian consumer preferences for beef using linked samples from Australian carcasses, (ii) between the actual scores of French and Australian beef given by French consumers (including medium and rare cooking) compared with those predicted using the MSA model.

Material and methods

Source of beef

Eighteen Australian cattle, all steers, were slaughtered at the Northern Co-operative Meat Company abattoir at Casino in

New South Wales (Australia). A part of this meat was dedicated to Australian consumer testing, another part to French consumer testing and a third part to beef testing in South Africa under another project (Thompson *et al.*, 2010).

Eighteen French cattle were slaughtered under EU requirements following French industry practice, in a private abattoir in the western part of France.

The type of cattle was chosen to reflect common commercial production offered to French consumers, which is very different from the Australian one. Three young bulls, three young dairy cows, six cull dairy cows and six cull beef cows were used to provide the French beef samples in the French experiment, none of them were steers (Table 1). The carcasses were weighed and graded according to the EU classification including conformation and fatness. In addition, pH/temperature declines on all the carcasses were recorded according to Perry *et al.* (2001). Moreover, the carcasses were graded by a qualified MSA grader for ossification, marbling score, meat colour, rib fat, ultimate pH and muscle temperature (Thompson, 2002).

Muscles

Six muscles were tested as paired samples in France and Australia. They were chosen to give a wide range of eating qualities: outside (*m. biceps femoris*), topside (*m. semimembranosus*), striploin (*m. longissimus dorsi*), rump (*m. gluteus medius*), oyster blade (*m. infraspinatus*) and tenderloin (*m. psoas major*).

For Australia, each muscle was collected from 18 animals with 9 of each muscle aged 5 days and 9 aged 21 days, whereas there was a single ageing time of 10 days for the French samples.

Meat preparation and French consumer panels

Consumer assessment of eating quality was done according to protocols for MSA testing described by Watson *et al.* (2008a).

Grilled steaks were cooked on a Silex clamshell grill (Silex, Hamburg, Germany) set to 220°C for 4.75 min for 'medium' and to 200°C for 3.25 min for 'rare' (Watson *et al.*, 2008a, Accessory publication).

Each consumer (Australian or French) received seven samples: the first sample was a link sample derived from either the striploin or rump eye muscles. Following this first

link sample, every consumer received a further six steaks, one from each of six different muscle types. The order of product presentation to consumers was controlled by a 6×6 Latin square, which ensured that each product was served an equal number of times in each presentational order position (two to seven) and an equal number of times before and after each other product to balance out any potential order, lag or halo effects. Within each of the products, there were six source samples (one muscle from one animal) comprising three from France and three from Australia. Ten consumers tasted each. This involved five steaks from each sample, which were halved after cooking. The five steaks from each sample were cooked in different rounds, so every sample was always spread across five of the six possible order positions from two to seven and in addition controlled so that one steak was served within five separate subgroups of 12 consumers within the base 60 consumers used in any one session.

Thus, in total, 360 French consumers took part in the 'medium' cooking test, with each consumer eating three Australian beef samples and three French beef samples. In addition to that, 180 French consumers tested the 'rare' beef, only from French origin and prepared as paired samples from the same cuts used for the 'medium' cooking. In Australia, 180 consumers were used to test the Australian beef steaks, which were all cooked to medium.

Consumers scored portions for tenderness (tn), juiciness (ju), flavour liking (fl) and overall liking (ov), and for this purpose they made a mark on a 100 mm line scale. In addition, they were asked to assign a quality rating to each sample: 'unsatisfactory', 'good everyday' (equivalent to 3*), 'better than everyday quality' (equivalent to 4*) or 'premium quality' (equivalent to 5*). Before the taste panel, data were obtained on the demographic profile of the consumers and their families, the families' meat purchase habits and their frequency and habits of beef consumption. After completion of the beef tasting, consumers were asked to quantify their willingness to pay for the different grades (unsatisfactory, good everyday, better than everyday and premium) of meat.

Data analysis

The relationship of the satisfaction grade to the four sensory scores was determined using discriminant analysis (Watson *et al.*, 2008a). The problem addressed here was to combine consumer scores for tn, ju, fl and ov into a single score (called MQ4) that could be used to predict the four satisfaction grades (namely 'unsatisfactory', 'good everyday', 'better than everyday quality' or 'premium quality'). A linear discriminant analysis was used with the satisfaction grade as the criterion to be predicted by tn, ju, fl and ov.

Results and discussion

Consumer characteristics

In total, 540 French consumers were involved, including 306 women (57%) and 234 men (43%). Overall, there was an even distribution between the six age classes, except for the

oldest people (>65 years) who were very few (3.3% only). Women were most represented in the young classes (<35 years) and men in the oldest one (>65 years). All the Australian consumers were aged between 20 and 50 years, with 54% in the 40- to 50-year age category. The gender ratio was 47% and 53% males and females, respectively.

Quality and meat price

The actual values for the willingness to pay were 5.0€ for unsatisfactory, 11.6€ for 3*, 16.2€ for 4* and 21.7€ for 5* (which means a 1.87-fold difference between 3* and 5* products). Differences in France were smaller than those observed in other countries such as Japan or the United States of America (with 2.94- and 2.42-fold differences between 3* and 5* products) but very similar to those from Ireland where a 1.64-fold difference was observed between 3* and 5* beef (Lyford *et al.*, 2010).

MQ4 as a measure of eating quality for French consumers

Values of consumer scores for tn, ju, fl and ov are shown in Table 2 for each group of French and Australian consumers. There was generally a good agreement between the palatability scores for French and Australian consumers eating the same samples at the same degree of doneness with an R^2 for tn, ju, fl and ov of 0.96, 0.79, 0.94 and 0.94, respectively. There was also a high correlation between medium and rare cooking for the French consumer palatability scores ($R^2 > 0.9$).

The optimal weightings of the four sensory parameters (tn, ju, fl and ov) to predict the final rating ('unsatisfactory', 3*, 4* and 5*) from the MQ4 score (equal to $0.31 \text{ tn} + 0.04 \text{ ju} + 0.30 \text{ fl} + 0.36 \text{ ov}$ and indicated in Table 2) were similar to those in Australia, as the weighting for tn, ju, fl and ov scores currently used for MSA in Australia are 0.3, 0.1, 0.3 and 0.3 (Thompson *et al.*, 2010). Moreover, these weightings predicted the actual rating given by consumers for over 70% of the total number of samples, indicating that a high level of prediction is possible for French consumers. The MSA boundaries for French consumers between 'unsatisfactory', 3*, 4* and 5* were found to be ca. 38, 61 and 80, respectively. The differences between extreme classes are therefore slightly more important in France than in Australia (Figure 1).

Prediction of beef quality

The Australian MSA model provides a predicted MQ4 score (MQ4pred) for each cut from the factors recorded for each carcass. One important area of interest was the differences between (i) the observed MQ4 score (MQ4obs) based on real consumer scores and calculated from the above equation ($\text{MQ4} = 0.31 \text{ tn} + 0.04 \text{ ju} + 0.30 \text{ fl} + 0.36 \text{ ov}$) and (ii) the MQ4pred from the Australian model. The mean differences (or residuals between observed and predicted values) are presented for each cut eaten by French consumers in Figures 2 and 3.

These results showed that, on average across all muscles consumed in France, the MSA model predicted the MQ4 score, with global predictive scores per country origin \times ageing time never deviating by more than 5 on a 0 to 100

Table 2 Means (\pm s.e.) for sensory scores given by Australian and French consumers according to the country origin of samples, and according to ageing time or cooking method depending on the groups of samples

Consumers, meat origin and number of samples	Ageing time (days)	Cooking doneness	Tenderness	Juiciness	Flavour	Overall liking	Observed MQ4
Australian consumers and Australian samples ($n = 18$)							
Outside	5	Medium	31 \pm 2.6	42 \pm 3.1	42 \pm 2.6	40 \pm 2.8	38 \pm 2.4
Outside	21		28 \pm 3.7	39 \pm 4.2	40 \pm 2.8	36 \pm 3.6	35 \pm 3.3
Oyster blade	5		64 \pm 3.7	71 \pm 2.4	65 \pm 3.1	66 \pm 2.8	65 \pm 3.0
Oyster blade	21		70 \pm 2.0	75 \pm 2.1	71 \pm 2.1	71 \pm 1.9	71 \pm 1.9
Rump	5		45 \pm 1.9	42 \pm 2.9	49 \pm 2.2	48 \pm 2.3	47 \pm 2.0
Rump	21		57 \pm 3.2	54 \pm 3.3	59 \pm 2.8	59 \pm 3.3	58 \pm 3.0
Striploin	5		58 \pm 6.4	57 \pm 4.6	56 \pm 5.5	57 \pm 5.6	57 \pm 5.5
Striploin	21		58 \pm 3.6	56 \pm 3.4	59 \pm 3.7	59 \pm 3.9	58 \pm 3.6
Tenderloin	5		78 \pm 2.5	71 \pm 3.7	74 \pm 4.1	76 \pm 3.6	76 \pm 3.3
Tenderloin	21		77 \pm 1.7	72 \pm 2.5	74 \pm 2.2	76 \pm 1.7	75 \pm 1.7
Topside	5		29 \pm 4.1	32 \pm 3.8	40 \pm 4.1	35 \pm 3.7	35 \pm 3.9
Topside	21		37 \pm 3.3	39 \pm 3.0	44 \pm 3.0	42 \pm 3.0	41 \pm 2.9
French consumers and Australia samples ($n=8$ to 10)							
Outside	5	Medium	37 \pm 3.1	44 \pm 2.6	48 \pm 3.2	43 \pm 2.9	43 \pm 2.7
Outside	21		33 \pm 3.8	41 \pm 3.4	43 \pm 3.5	38 \pm 3.5	38 \pm 3.5
Oyster blade	5		78 \pm 1.9	73 \pm 2.7	74 \pm 2.4	76 \pm 2.3	76 \pm 2.1
Oyster blade	21		79 \pm 1.9	67 \pm 2.2	72 \pm 2.0	74 \pm 2.2	74 \pm 1.9
Rump	5		58 \pm 4.2	53 \pm 2.8	59 \pm 2.7	57 \pm 3.4	57 \pm 3.3
Rump	21		57 \pm 2.6	47 \pm 2.8	56 \pm 1.9	54 \pm 2.0	55 \pm 2.1
Striploin	5		64 \pm 3.8	53 \pm 4.2	61 \pm 3.3	60 \pm 3.7	61 \pm 3.5
Striploin	21		69 \pm 2.7	51 \pm 3.6	61 \pm 2.2	62 \pm 2.4	63 \pm 2.4
Tenderloin	5		86 \pm 2.2	60 \pm 3.9	78 \pm 2.8	80 \pm 2.3	79 \pm 2.4
Tenderloin	21		84 \pm 1.6	66 \pm 3.7	77 \pm 1.8	77 \pm 1.8	78 \pm 1.6
Topside	5		32 \pm 4.4	36 \pm 3.0	43 \pm 3.5	37 \pm 4.1	37 \pm 3.8
Topside	21		33 \pm 2.6	33 \pm 2.5	41 \pm 2.7	36 \pm 2.6	36 \pm 2.5
French consumers and French samples ($n=18$)							
Outside	10	Medium	24 \pm 2.6	42 \pm 2.5	40 \pm 2.1	32 \pm 2.3	33 \pm 2.1
Outside	10	Rare	25 \pm 2.3	44 \pm 1.7	42 \pm 2.5	32 \pm 2.3	34 \pm 2.2
Oyster blade	10	Medium	58 \pm 3.0	64 \pm 2.6	58 \pm 2.4	58 \pm 2.5	59 \pm 2.4
Oyster blade	10	Rare	68 \pm 2.4	69 \pm 1.4	62 \pm 2.3	62 \pm 2.3	64 \pm 2.0
Rump	10	Medium	54 \pm 3.5	53 \pm 3.4	57 \pm 2.6	56 \pm 2.9	55 \pm 3.0
Rump	10	Rare	56 \pm 2.3	62 \pm 2.0	59 \pm 2.4	57 \pm 2.4	58 \pm 2.2
Striploin	10	Medium	51 \pm 3.6	54 \pm 2.5	56 \pm 1.9	54 \pm 2.7	54 \pm 2.6
Striploin	10	Rare	62 \pm 2.3	63 \pm 2.2	62 \pm 2.2	62 \pm 2.2	62 \pm 2.0
Tenderloin	10	Medium	83 \pm 1.5	68 \pm 2.8	76 \pm 1.6	78 \pm 1.5	78 \pm 1.5
Tenderloin	10	Rare	87 \pm 1.3	71 \pm 2.7	77 \pm 2.2	78 \pm 2.2	80 \pm 1.8
Topside	10	Medium	36 \pm 3.3	42 \pm 2.8	47 \pm 2.1	41 \pm 2.5	42 \pm 2.5
Topside	10	Rare	43 \pm 3.0	51 \pm 2.1	53 \pm 2.4	48 \pm 2.6	49 \pm 2.5

Each sample was tested by 10 consumers. Each sample score was the mean of the assessment by these 10 consumers. The table indicates means \pm s.e. of sample scores for nine Australian samples at each ageing time, and for 18 French samples for each cooking method.

scale (Figure 2). The quality score was rather underestimated or overestimated about 2 points for all samples. However, there were three main discrepancies, regarding the results for each individual muscle. The oyster blade was undervalued by at least 5 points for the Australian samples regardless of ageing time, but overvalued by almost 6 points for the French samples. French topside and rump were also underevaluated by 8 to 12 points, but not the Australian ones. Only the outside was clearly overvalued by 7 to 10 points for the samples aged

more than 5 days (French 10 days and Australian 21 days): the ageing effectiveness was obviously overestimated for this rather tough muscle (Figure 3).

It is also interesting to consider the results by muscle and animal type, given that there were different confounding effects (Figure 3). Muscle samples from steers were aged either 5 or 21 days, and young bull and cow samples were aged 10 days. Meanwhile, muscles from steers were only tested at a 'medium' doneness, whereas for young bulls and




Figure 1 Meat Standards Australia boundaries between 'unsatisfactory', 3*, 4* and 5* in France and in Australia.




Figure 2 Mean differences (+/- standard errors) between MQobs and MQpred for muscles tested by French consumers at different ageing times and also global average across all steaks (all).




Figure 3 Mean differences (+/- standard errors) between MQobs and MQpred for meats tested by French consumers, by animal types and muscles.

cows half were consumed 'medium' and half were consumed 'rare'. The main discrepancy with Australian steers was that the oyster blade was slightly undervalued. There were also small discrepancies with the outside, the topside and the rump muscles from young French bulls. Apart from that, French meat from young French bulls was well predicted by the steer prediction from the Australian MSA system, which at this stage does not have a prediction for meat from bulls. On the other hand, quality of French cow beef seemed to be mainly undervalued by the MSA prediction model except for the outside and the oyster blade (Figure 3).

Future perspectives

This study indicated that the French beef quality can be predicted using the current MSA system as for the Australian beef. This conclusion was also found in previous studies in Japan (Polkinghorne RJ, personal communication), Korea

(Thompson *et al.*, 2008) and the United States of America (Smith *et al.*, 2008).

One of the goals of the current European research programme called ProSafeBeef (2007 to 2011), in which France, Ireland, Germany, Belgium, Greece, the United Kingdom (among the major partners) are participating, is to try to establish a European prediction model for beef. This could draw on the Australian MSA system for inspiration and expertise. Another goal of the ProSafeBeef programme is to include in this meat quality prediction model biochemical characteristics of muscles related to meat quality (Hocquette *et al.*, 2011b) and the genomic markers that have been discovered in various countries (Hocquette *et al.*, 2007; Hocquette *et al.*, 2010). The MSA system is indeed currently considering the incorporation of gene markers for tenderness into the model.

Another important question in France would be to test beef from different pure breeds to those commonly used in Australia. Thus, common French cattle breeds such as Limousine and Charolais are typically used for cross-breeding in Australia, and given the typically lower levels of intramuscular fat and different muscle structure (Jurie *et al.*, 2007) further work is required to predict the eating quality for these cattle as pure breeds. This is important from a French perspective in order to support existing official quality signs. In fact, the MSA system is a meticulous approach capable of supporting the pre-existing quality signs without aiming to enter into competition with them.

Comparison of consumer data across countries with the same beef samples would also be essential to start the development of a European system to predict quality. Indeed, recent work undertaken within the European research programme ProSafeBeef showed that the concept of a beef eating-quality guarantee system is well accepted by European beef consumers (Verbeke *et al.*, 2010).

Conclusion

The MSA system has been proved to be able to predict beef quality in many countries throughout the world, suggesting common attitudes of consumers regardless of the country and despite culture differences across them (for instance, for cooking beef). Nevertheless, further studies are still necessary to report the effectiveness of the MSA model over a wider range of treatments.

The MSA system represents the first elaborate system to predict the sensorial guarantee of a cut of beef according to its length of ageing and the method of cooking. The system must be praised for having created a federation in Australia with a large number of beef industry personnel involved, including scientists, farmers, processors and retailers. This trans-chain approach aimed at satisfying consumer expectations immediately raises questions concerning traditional attitudes and the positioning of the different professionals in the beef chain, at least in France. The French beef industry still needs to be convinced and to express its wishes.

Acknowledgements

The authors thank the financial contributions of the 'National Office for Meat and Dairy Products' (now 'FranceAgriMer'), of the French 'Direction Générale de l'Alimentation' and of a French well-known private Company, which also provided the meat of 3 of the 18 animals needed for the experiment. Another small private enterprise specialized in meat selection and boning used for gourmet restaurants (Jean Denaud) was also associated to this experiment. Two organizations dealing with the Charolais breed ('Association Institut du Charolais', 'Syndicat de Défense et de promotion de la Viande de Boeuf de Charolles') also funded this experiment. Apart from that, French and Australian journeys were supported for 2 years by 'Egide/Fast' funds from French and Australian governments, respectively.

The authors would like to acknowledge the assistance of Janine Lau and Alan Gee (Australia). The authors also thank John Thompson (Australia) and Paul Allen (Ireland) for their help.

References

- Ferguson D, Thompson J and Polkinghorne R 1999. Meat Standards Australia, a "PACCP" based beef grading scheme for consumers. 3) PACCP requirements which apply to carcass processing. Book of Abstracts of the 45th International Congress of Meat Science and Technology, Yokohama, Japan, pp. 18–19.
- Hocquette JF and Chatellier V 2011. Prospects for the European beef sector over the next 30 years. *Animal Frontiers* 1, 20–28.
- Hocquette JF, Gondret F, Baéza E, Médale F, Jurie C and Pethick DW 2010. Intramuscular fat content in meat-producing animals: development, genetic and nutritional control, identification of putative markers. *Animal* 4, 303–319.
- Hocquette JF, Legrand I, Jurie C, Pethick D and Micol D 2011a. Perception of the Australian system for the prediction of beef quality (MSA) with perspectives for the European beef sector. *Animal Production Science* 51, 30–36.
- Hocquette JF, Lehnert S, Barendse W, Cassar-Malek I and Picard B 2007. Recent advances in cattle functional genomics and their application to beef quality. *Animal* 1, 159–173.
- Hocquette JF, Meurice P, Brun JP, Jurie C, Denoyelle C, Bauchart D, Renand G, Nute GR and Picard B 2011b. The challenge and limitations of combining data: a case study examining the relationship between intramuscular fat content and flavour intensity based on the BIF-BEEF database. *Animal Production Science* 51, 975–981.
- Hwang IH, Polkinghorne R, Lee JM and Thompson JM 2008. Demographic and design effects on beef sensory scores given by Korean and Australian consumers. *Australian Journal of Experimental Agriculture* 48, 1387–1395.
- Jurie C, Cassar-Malek I, Bonnet M, Leroux C, Bauchart D, Boulesteix P, Pethick DW and Hocquette JF 2007. Adipocyte fatty acid-binding protein and mitochondrial enzyme activities in muscles as relevant indicators of marbling in cattle. *Journal of Animal Science* 85, 2660–2669.
- Lyford C, Thompson JT, Polkinghorne RJ, Miller MF, Nishimura T, Neath K, Allen P and Belasco E 2010. Is willingness to pay (WTP) for beef quality grades affected by consumer demographics and meat consumption preferences? *Australasian Agribusiness Review* 18, 1–17.
- Moëvi I, Hocquette JF, Jurie C and Micol D 2008a. Expertise du système australien de prédiction de la qualité de la viande bovine (MSA). Quelles perspectives pour la filière française ? *Compte rendu n° 170832010 (TEND-06-058)*. Institut de l'Elevage, Inra, Interbev, Office de l'Elevage, Paris, France.
- Moëvi I, Hocquette JF, Jurie C and Micol D 2008b. The Australian beef eating quality prediction model MSA: presentation, scientific and professional assessment. *Rencontres autour des Recherches sur les Ruminants* 15, 97–100.
- Perry D, Shorthose WR, Ferguson DM and Thompson JM 2001. Methods used in the CRC program for the determination of carcass yield and beef quality. *Australian Journal of Experimental Agriculture* 41, 953–957.
- Polkinghorne R, Watson R, Porter M, Gee A, Scott J and Thompson J 1999. Meat Standards Australia, a "PACCP" based beef grading scheme for consumers. 1) The use of consumer scores to set grade standards. Book of Abstracts of the 45th International Congress of Meat Science and Technology, Yokohama, Japan, pp. 14–15.
- Smith GC, Tatum JD and Belk KE 2008. International perspective: characterization of United States Department of Agriculture and Meat Standards Australia systems for assessing beef quality. *Australian Journal of Experimental Agriculture* 48, 1465–1480.
- Thompson JM 2002. Managing meat tenderness. *Meat Science* 60, 365–369.
- Thompson J, Polkinghorne R, Hearnshaw H and Ferguson D 1999a. Meat Standards Australia, a "PACCP" based beef grading scheme for consumers. 2) PACCP requirements which apply to the production sector. Book of Abstracts of the 45th International Congress of Meat Science and Technology, Yokohama, Japan, pp. 16–17.
- Thompson J, Polkinghorne R, Watson R, Gee A and Murison B 1999b. Meat Standards Australia, a "PACCP" based beef grading scheme for consumers. 4) A cut based grading scheme to predict eating quality by cooking method. Book of Abstracts of the 45th International Congress of Meat Science and Technology, Yokohama, Japan, pp. 20–21.
- Thompson JM, Polkinghorne R, Hwang IH, Gee AM, Cho SH, Park BY and Lee JM 2008. Beef quality grades as determined by Korean and Australian consumers. *Australian Journal of Experimental Agriculture* 48, 1380–1386.
- Thompson JM, Polkinghorne RJ, Gee A, Motiang D, Strydom P, Mashau M, Ng'ambi J, de Kock R and Burrow H 2010. Beef palatability in the Republic of South Africa: implications for niche-marketing strategies. *ACIAR Technical Reports* 72. Australian Centre for International Agricultural Research (ACIAR), Canberra, Australia.
- Verbeke W, Wezemael LV, de Barcellos MD, Kügler JO, Hocquette JF, Ueland O and Grunert KG 2010. European beef consumers' interest in a beef eating-quality guarantee: insights from a qualitative study in four EU countries. *Appetite* 54, 289–296.
- Watson R, Gee A, Polkinghorne R and Porter M 2008a. Consumer assessment of eating quality – development of protocols for Meat Standards Australia (MSA) testing. *Australian Journal of Experimental Agriculture* 48, 1360–1367. Accessory publication: http://www.publish.csiro.au/?act=view_file&file_id=EA07176_AC.pdf
- Watson R, Polkinghorne R and Thompson JM 2008b. Development of the Meat Standards Australia (MSA) prediction model for beef palatability. *Australian Journal of Experimental Agriculture* 48, 1368–1379.